

SHOT Report 2017

Errata list

This list is not guaranteed to be complete. Errors such as minor typing errors or missing superscript fonts are not listed. SHOT would be grateful if you would report any further major or serious errors if you discover them.

Page	Item / Location	Error	Notes / Corrected text
11	Figure 3.2	The number of preventable deaths should be 14 not 12, as the preventable deaths include the 2 deaths related to Under and overtransfusion.	The corrected figure is displayed below, and this has been corrected on all the online content.
135	Figure 17.1	The number of suspected viral incidents reported and investigated should be 8 not 7.	The corrected figure is displayed below, and this has been corrected on all the online content.
153	Commentary under Case 18b.2	There is an error in the calculation used for raising the Hb level. Based on 0.4mL/kg, the level should be 1g/L and not 10g/L. BSH guidelines state that for adults of low body weight, a calculation of 4 mL kg ⁻¹ of red blood cells in additive solution will produce an approximate rise of 10 g/L ⁻¹ (https://onlinelibrary.wiley.com/doi/full/10.1111/tme.12481)	Based upon a calculation of 0.4mL/kg raising the Hb level by 1g/L This has been corrected on all the online content.
Summary	Page 1	The number of preventable deaths should be 14 not 12	21 deaths, 14 preventable This is also correct on the online version of the summary

Figure 3.2: Deaths related to transfusion (with imputability) reported in 2017 n=21




Figure 17.1: Outcome of reports of suspected TTI made to the NHSBT/PHE Epidemiology Unit in 2017*

